

**INGREDIENTS:** Contains strains of *Lactobacillus acidophilus* and *Pediococcus acidilacticii*, skim milk, fructo-oligosaccharides and inactivated *Saccharomyces cerevisiae*.

**DIRECTIONS FOR USE:** Use of a quality water acidifier for at least 8 hours prior to administration of FloraMax-B11 will enhance delivery of beneficial microflora. Mix 1 bottle for up to 20,000 birds in stock solution sufficient to provide birds with product for 6-12 hours. Use of a vaccine stabilizer is recommended. Discard any remaining product after 24 hours. Disinfectant or antibiotic residues may kill the microflora in this product. Discontinue water sanitizers 24 hours prior to and 24 hours after administration.

**STORAGE:** Keep Refrigerated. Brief exposure to room temperatures will not harm the product.

**MANUFACTURED BY:** PACIFIC VET GROUP USA, INC., 2135 Creek View, Fayetteville, Arkansas, USA 72704  
©2008 Pacific Vet Group USA, Inc.



# FloraMax® -B11

**A Probiotic Formulation  
Specifically for Use in Poultry**

MANUFACTURED UNDER EXCLUSIVE  
LICENSE FROM THE UNIVERSITY OF ARKANSAS

KEEP REFRIGERATED

**NET CONTENT 140g (20,000 DOSES)**